



Contents lists available at ScienceDirect

Epilepsy & Behavior

journal homepage: www.elsevier.com/locate/yebeh

Review

Common psychiatric comorbidities in epilepsy: How big of a problem is it?

Jay A. Salpekar^{a,*}, Marco Mula^{b,c}^a Kennedy Krieger Institute, Johns Hopkins University, USA^b Institute of Medical and Biomedical Education, St George's University of London, United Kingdom^c Atkinson Morley Regional Neuroscience Centre, St George's University Hospitals NHS Foundation Trust, London, United Kingdom

ARTICLE INFO

Article history:

Received 23 July 2018

Accepted 24 July 2018

Available online xxx

Keywords:

Psychiatry

Epilepsy

Depression

Cognitive

Comorbidity

Behavior

Diagnosis

ABSTRACT

Psychiatric illness and epilepsy commonly co-occur in adults and in children and adolescents. Theories of comorbidity are complex, but recurring associations between the conditions suggest overlap that is more than simple co-occurrence. Common underlying pathophysiology may imply that epilepsy itself may constitutively include psychiatric symptoms. Conditions such as depression or cognitive difficulties commonly occur and in some cases, are considered to be associated with specific epilepsy characteristics such as localization or seizure type. Regardless of etiologic attributions to psychiatric comorbidity, it is clear today that treatment for epilepsy needs to target psychiatric illness. In many cases, quality-of-life improvements depend more upon addressing psychiatric symptoms than seizures themselves.

© 2018 Elsevier Inc. All rights reserved.

1. Introduction

Psychiatric comorbidities have long been studied in the context of epilepsy. A sizeable body of evidence supports the understanding that psychiatric illness is overrepresented in epilepsy as compared with other chronic medical illnesses [1]. Even with the markedly altered functionality and lifestyle associated with conditions such as cancer, diabetes, or asthma, the presence of psychiatric comorbidity in epilepsy appears to be more substantial [2]. Although opinions regarding the best methods of diagnostic assessment may differ, comparable assessment tools clearly find that the overrepresentation of psychiatric illness is significant both in adults and in children [3,4].

Nevertheless, a persisting challenge is in measuring the presence of comorbidity independent of possible confounding factors. Very few studies assessing comorbidity are prospective in nature or have a population-based approach. Additionally, the broad spectrum of illness makes it difficult to isolate psychiatric symptoms from underlying medical or neurologic illness. The population with epilepsy itself is heterogeneous, so overlap with heterogeneous and somewhat arbitrarily categorized psychiatric conditions may hinder a comprehensive diagnostic approach. Still, even with these challenges, the connection

between psychiatric illness and epilepsy is abundantly present, especially for mood disorders and problems of cognition.

What remains unclear is how epilepsy-related characteristics may be etiologically associated with psychiatric comorbidity. While it has been long considered intuitive that the more severe the epilepsy, the more severe the comorbidity, this may not be completely accurate. It may be that specific seizure-related characteristics affect the presence or absence of comorbidity. Seizure frequency is commonly viewed as adversely affecting psychiatric state, as is the usage of multiple anticonvulsant medications, thus, adding to the potential risk of developing psychiatric illness. However, neither context may represent the essence of comorbidity in the truest sense.

Ultimately, it may be that epilepsy is a broad spectrum illness with heterogeneity that includes psychiatric aspects. In that sense, thinking of comorbidity as the association of two etiologically unrelated conditions would be inaccurate. Certain types of epilepsies could have psychiatric, cognitive, or behavioral complications that are inherently associated with a common underlying pathophysiology. With that approach, comorbidity would not exist; instead, the associated behavioral symptoms would be viewed simply as constituent parts of the epilepsy.

Such a paradigm shift has marked implications in terms of treatment. Despite the intrigue in thinking of comorbidity in this manner, it is a modern approach, and as such, the evidence base for treatment is underdeveloped. Treatment studies and clinical trials are not typically conceptualized in this way. However, it seems clearer now that in order to comprehensively treat epilepsy and to improve quality of life, both

* Corresponding author at: Neuropsychiatry in Epilepsy Program, Kennedy Krieger Institute, Johns Hopkins University School of Medicine, 716 N. Broadway, Baltimore, MD 21205, USA.

E-mail address: jsalpek1@jhmi.edu (J.A. Salpekar).

behavioral and seizure specific outcomes need to be addressed. Even though this paradigm does not reflect the state of the evidence at this time, the approach of identifying psychiatric symptoms independently is still meaningful as a first step in management. The evidence base does help us in that regard and will be reviewed in the following sections.

2. Common comorbidities in adults

2.1. Epidemiological aspects in adults with epilepsy

2.1.1. Mood disorders

Cross-sectional, population-based studies in adults with epilepsy are showing a uniformly increased prevalence of depression, with prevalence rates very similar to those of the general population only among seizure-free patients [5], while the prevalence is definitely higher than that in unselected samples (between 17% and 22%) [6], and it is up to 55% in patients with drug-resistant epilepsy [7]. In general terms, these figures partially reflect the severity of the underlying seizure disorder not only in terms of psychosocial difficulties but also in terms of brain dysfunction. Cross-sectional studies are highly informative from a public health perspective, as they give an idea of the size of the problem, but cohort studies can provide information on the temporal relationship between comorbid conditions. In fact, a number of studies are now suggesting that the relationship between epilepsy and depression is not necessarily unidirectional, and patients with depression are at increased risk of developing epilepsy as well. Data from the UK General Practice Research Database show that the incidence-rate ratio of depression is significantly higher during the three years preceding the onset of epilepsy [8]. A study from Sweden shows that the age-adjusted odds ratio for the development of epilepsy is 2.5 for patients with a depressive disorder [9], and these figures have been confirmed by at least three other studies with an increased risk of up to 7 times in selected cases [10–12]. All these data clearly suggest that either some patients with depression develop epilepsy as part of the “natural course” of the depressive disorder or that depression is a premorbid phase of some epileptic syndromes.

2.1.2. Anxiety disorders

Data on anxiety disorders are less systematic than those on depression, and this is due to a number of reasons including the high prevalence of depression in epilepsy that can mask anxiety disorders, a common attitude among neurologists in considering anxiety as the natural consequence of having unpredictable seizures and the potential misdiagnosis between panic attacks and seizure-based phenomena like ictal fear. As already discussed in the context of depression, the few available studies suggest a uniformly increased prevalence of all anxiety disorders [13]. Two US National surveys show that people with self-reported epilepsy are two times more likely to report a diagnosis of anxiety disorder than those without [14,15]. These figures have been replicated by another cross-sectional, population-based study in unselected patients with epilepsy using standardized clinical interviews [16]. Although less established as compared with depression, a few preliminary studies are suggesting a similar bidirectional relationship between epilepsy and anxiety disorders. A US study in veterans older than 65 shows that a previous history of anxiety is significantly more common in those who developed epilepsy as compared with controls [17]. A population-based, case-control study in Sweden shows that patients hospitalized for anxiety disorders are more than 2 times at increased risk of developing unprovoked seizures [9]. Finally, a cohort study using data from the UK General Practice Research Database shows that the incidence for anxiety disorders is not only higher in people with epilepsy as compared with controls but is also already increased three years prior to the diagnosis of epilepsy [8].

2.1.3. Psychoses

Psychoses and thought disorders are less frequent than mood and anxiety disorders, but they represent serious complications affecting prognosis, morbidity, and mortality in epilepsy. In general terms, epidemiological studies show that the prevalence of psychoses is 7 times higher than that of schizophreniform disorders in the general population [18]. Psychotic symptoms in epilepsy are well recognized in a number of different clinical scenarios, from postictal psychoses [19] to antiepileptic drug-related psychotic episodes [20] to chronic schizophrenia-like disorders [21]. The prevalence of psychoses in epilepsy seems again related to the severity of the seizure disorder but also to the degree of pathology of the temporolimbic structures. Cross-sectional, population-based studies in unselected samples show that the prevalence of nonorganic, nonaffective psychoses, including schizophrenia and related disorders, is around 4%–5% [9,22] but in selected samples, such as hospital case series, is even higher than that [23]. A systematic review and meta-analysis of a published literature pointed out that patients with temporal lobe epilepsy present with the highest prevalence in the range of 7% [24]. In patients with temporal lobe epilepsy, those with febrile seizures and hippocampal sclerosis are more likely to develop psychoses than those with normal or minimal changes on Magnetic Resonance Imaging (MRI) [25]. Interestingly, both a family history of psychoses and a family history of epilepsy have been identified as potential risk factors suggesting a strong neurobiological connection between epilepsy and psychoses [18]. As already discussed in the context of depression, psychoses have also shown a bidirectional relationship with epilepsy. A case-control study in Sweden shows that the age-adjusted odds ratio for seizures is 2.5 in patients with any diagnosis of psychosis even after controlling for antipsychotic drug treatment and other potential confounding factors [9]. Two retrospective cohort studies show that patients with schizophrenia have 2 to 3 times increased risk to develop epilepsy [26] with incidence rates of 7 per 1000 person-year [27].

2.1.4. Cognitive problems

Cognitive problems represent a significant burden for any neurological disorder. Despite the large amount of epidemiological studies in children with epilepsy, there are no population-based studies investigating the neuropsychological status of people with adult-onset epilepsy [28]. This obviously represents a serious omission in the scientific literature as the issue of cognitive problems is not just for children and young people but also for adults. In fact, a subjective impairment in cognitive functions is frequently reported by adult patients and ranging between 44% and 59% [29]. Most importantly, 63% of patients perceive that antiepileptic drugs prevent them from achieving goals in life rather than the epilepsy itself [30]. The problem of neuropsychological deficits in adult-onset epilepsies is quite complex. Several studies have clearly shown that subjective memory complaints do not correlate with formal neuropsychological deficits but are rather due to other factors, especially anxiety and depression [31,32]. Antiepileptic drugs have a different propensity for cognitive side effects, but a number of different variables, which are both patient-related and epilepsy-related, have to be taken into account in adults [30].

Patients are often concerned that epilepsy may be responsible for accelerated brain aging. Although population-based studies in this area are lacking, it has been suggested that the potential cognitive decline in aging patients with epilepsy is very slow, and the cognitive performance at the onset of the epilepsy is the most important variable [28, 33]. It has been suggested that people with epilepsy may reach a clinically significant threshold of impairment late in life because they start at a lower baseline cognitive level than healthy subjects rather than because of a true accelerated cognitive decline [33]. This would be further supported by studies in newly diagnosed adult patients with epilepsy showing that they are cognitively compromised even at the onset of the disorder and before starting antiepileptic drug treatment [34]. Further studies in this area are urgently needed.

2.2. Psychiatric comorbidities as a poor prognostic marker in adults with epilepsy

The amount of epidemiological data on psychiatric comorbidities of epilepsy has increased the attention of clinicians and researchers on the potential impact of comorbidities on the prognosis of epilepsy and the identification of potential biomarkers [35]. As already discussed, the different prevalence rates in different populations with epilepsy, namely seizure-free patients and patients with drug-resistant epilepsy, seem to suggest that the presence of psychiatric comorbidities partially reflects the severity of the seizure disorder. In adults with epilepsy, there is now enough literature supporting the hypothesis that psychiatric comorbidities represent also a poor prognostic marker.

It has been established a long time ago that depression is a better indicator of quality of life than seizure frequency itself [36]. It is now becoming evident that depression is also associated with poor response to antiepileptic drug treatment [37,38], poor outcome after epilepsy surgery [39,40], increased seizure severity [41], increased risk of injury [42], and premature mortality [43]. Future studies need to clarify whether early identification and prompt treatment of psychiatric comorbidities can have an impact on the prognosis of the epilepsy or whether they just represent indicators of poor prognosis.

3. Comorbidities in pediatrics

Psychiatric comorbidity has historically been less well studied in children than in adults. However, recent high-quality population studies offer detailed information regarding the amount of medical and psychiatric comorbidity in pediatric epilepsy. A recent study of a Norwegian patient registry showed that 78.3% of the pediatric epilepsy population had comorbid medical, neurologic, or psychiatric illness as compared with 30.3% of the general pediatric population [44]. Developmental or psychiatric disorders were present in 42.9% of children with epilepsy and higher if the epilepsy was deemed complicated. Although the diagnoses were not verifiable beyond the initial medical record notation, the sample size of over one million is notable and reinforces the notion that psychiatric comorbidity is significant in the larger population of persons with epilepsy.

Sizeable comorbidity has also been found in a large-scale retrospective study of newly diagnosed epilepsy. Administrative review of over six million records found 7654 children with newly diagnosed epilepsy. Neurobehavioral comorbidities were present in 60% of these children as compared with 23% without epilepsy [45]. These recent studies complement existing information in the adult population and affirm that comorbidity is markedly present in children as well as in adults. Furthermore, in children, the comorbidity may be more readily measurable and thus, stimulate valuable insight into theories of pathophysiologic overlap.

3.1. Behavioral side effects of anticonvulsants

A common question, especially in children, is whether anticonvulsants cause behavioral or cognitive side effects. Cognitive side effects may be more noticeable in children because of academic emphases and measurable trajectories of cognitive development. Much literature in epilepsy has been devoted to potential side effects of anticonvulsant drugs, and despite the difficulty in isolating etiologies, some medicines have been nevertheless identified as more likely associated with behavioral issues [46].

Behavioral side effects may be reported with any anticonvulsant, but depression and fatigue have commonly been associated with phenobarbital [47]. A recent retrospective review suggests that levetiracetam and valproate have behavioral side effects more often than other anticonvulsants, particularly if hyperactivity is noted at baseline [48].

It must also be noted that anticonvulsant drugs often have positive effects upon behavior, particularly in the context of mood disorders, and have proven effective either as primary or adjunct treatments [49]. The conundrum of anticonvulsants yielding behavioral or cognitive

side effects while at the same time, serving as a valuable treatment option is still present.

3.2. Childhood absence epilepsy (CAE)

Childhood absence epilepsy is characterized by interruptions in consciousness though with little other visible semiology. Inattention is obvious not only during seizures themselves but also has long been observed in these children in between absence episodes. One study of CAE comorbidity found that 25% had subtle cognitive deficits, 43% linguistic difficulties, and 61% had a psychiatric illness, primarily Attention-Deficit Hyperactivity Disorder (ADHD) [50]. Differentiation of CAE from the inattentive subtype of ADHD may be difficult [51].

For many years, CAE was considered a benign form of epilepsy, having little sequelae beyond the distraction associated with the discrete events. However, if this was the case, then comorbidities of attention or executive function deficits in children with epilepsy would not be markedly overrepresented as compared with the typical pediatric population. Instead, the cognitive deficits associated with CAE may be similar to those associated with ADHD in children without epilepsy [52]. Childhood absence epilepsy has increasingly become viewed as a disease with persisting cognitive problems, well beyond the time scale of individual absence episodes [53].

The idea that attention and cognitive problems are so prominent in CAE suggests physiologic overlap of cognition and the mechanisms that underlie absence seizures. An intriguing study recently done with functional magnetic resonance imaging (fMRI) suggests that connectivity in attention, salience, and default mode networks is altered in CAE [54]. The consideration is that these network abnormalities persist beyond discrete seizure episodes and may explain the problems of executive function commonly seen in CAE. Network abnormalities in epilepsy as well as in psychiatric disorders in general may explain a great deal of the overlap not only with CAE but also with other epilepsies. Childhood absence epilepsy is not just associated with attention or cognitive problems but may involve mood complications as well [55]. The widespread networks subsuming generalized epilepsy may intuitively explain comorbidities present, even in a pediatric population [56].

3.3. Mood disorder and anxiety

Mood disorder and anxiety often co-occur in children and adolescents without epilepsy. Co-occurrence of mood and anxiety disorders and epilepsy is likely to be as common in children as it is in adults, although large-scale prospective studies addressing this comorbidity have not been done [57]. Some studies in selected populations suggest co-occurrences of 5–40%, though samples are heterogeneous and assessment tools are variable [58–61]. Depression is underreported in children in general, and the same may be true in pediatric epilepsy [62,63]. Many with depressive symptoms do not seek mental health treatment for reasons of stigma or lack of clinician availability [64]. Clinically, quality-of-life measures have highlighted the adverse effect of depression in pediatric epilepsy. Similar to adults, the presence of depression has a marked effect upon quality of life and upon recovery postsurgery [65,66].

The association of depression or anxiety with specific seizure types or localization is less well established. However, two small studies suggest that depression is more often present in children with temporal lobe seizure localization [67,68]. This finding is consistent with literature associating depressive phenomena with temporal lobe pathology in persons without epilepsy [69].

3.4. ADHD

Attention-deficit hyperactivity disorder has proven to be very common in children with epilepsy. Reports estimate that 30% of children with epilepsy may meet criteria for ADHD, with some reports

suggesting prevalence up to 80% [70–72]. Both epilepsy and ADHD have problematic effects upon quality of life. A recent report suggests that each condition is significantly detrimental to quality of life, and in combination, the adverse effects upon quality of life may be additive [73].

It is still unclear as to which epilepsy characteristics are related to the presence of ADHD. One preliminary report suggests that attention problems are more common with localization-related epilepsy than with primarily generalized epilepsy, but the sample was somewhat skewed, and psychiatric diagnoses were not obtained at baseline [74]. Another study found that 80% of children with frontal lobe epilepsy with sustained abnormal discharges on electroencephalogram (EEG) also met criteria for ADHD [75].

Several reports suggest that EEG abnormalities may be present in children with ADHD even without confirmed epilepsy [76,77]. One report found that treatment with valproate improved paroxysmal EEG abnormalities as well as ADHD ratings [78]. The overlap may not be coincidental as a recent population study suggests that a bidirectional relationship is present between the two conditions. The presence of either ADHD or epilepsy increased the odds for the subsequent development of the other condition [79].

3.5. Autism spectrum disorders

The overlap between epilepsy and intellectual and developmental disabilities has been frequently reported. Epilepsy may be present in half of children and adolescents with autism or other developmental disorders [80]. Family studies also show overrepresentations of EEG abnormalities in persons with family histories of both autism and epilepsy [81]. Autism spectrum disorders are characterized by atypical communication, deficits in social reciprocity, and stereotyped or narrow ranges of behaviors or activities. Some epilepsy syndromes, such as Landau-Kleffner syndrome, include irregular communication styles or atypical social pragmatic language, similar to what is present in autism [82].

Ultimately, the overlap may be physiologic in nature. Some evidence suggests that neuronal hyperexcitability may be present in autism spectrum disorders. Epileptiform abnormalities have been reported in autism, in the absence of clinically evident seizures [83]. Such neuronal hyperexcitability would be expected in seizure disorders, and its presence in autism could reflect common pathophysiology between the two conditions. One study suggests that a deficit in interneurons mediating the inhibitory neurotransmitter gamma-aminobutyric acid (GABA) may be at fault in autism [84]. Such deficits in inhibition would be intuitive in epilepsy, which is in one sense, a disorder involving dysregulated neurons that activate inappropriately [85].

4. Conclusion

The weight of the evidence, regardless of heterogeneity in sampling or disagreements about ideal assessment techniques, clearly affirms that psychiatric comorbidity is a significant problem. Details about bidirectionality, etiology, or the possible existence of common pathophysiology are important, but the most important aspect may be the fact that these debates exist at all. The discourse itself results in part from the still underappreciated notion that psychiatric conditions are associated with epilepsy more often than expected.

Ultimately, medical care and treatment decisions relate to quality of life. Quality of life may necessitate bold approaches to treatment. Neurologists are the key directors of epilepsy care and as such, have no choice but to become competent in knowledge of psychiatry. For persons with epilepsy, psychiatric comorbidity appears to be the rule rather than the exception, and as such, comprehensive care demands attention to mental health.

Declarations of interest

JS (Lundbeck: Research funding to institution only); MM: none.

References

- Josephson CB, Jette N. Psychiatric comorbidities in epilepsy. *Int Rev Psychiatry* 2017; 29:409–24.
- Hermann BP, Vickrey B, Hays RD, Cramer J, Devinsky O, Meador K, et al. A comparison of health-related quality of life in patients with epilepsy, diabetes and multiple sclerosis. *Epilepsy Res* 1996;25:113–8.
- Jones JE, Siddarth P, Almame D, Gurbani S, Hermann BP, Caplan R. Identification of risk for severe psychiatric comorbidity in pediatric epilepsy. *Epilepsia* 2016;57: 1817–25.
- Hesdorffer DC, Baldin E, Caplan R, Berg AT. How do we measure psychiatric diagnoses? Implications of the choice of instruments in epilepsy. *Epilepsy Behav* 2014;31: 351–5.
- Jacoby A, Baker GA, Steen N, Potts PWCD. The clinical course of epilepsy and its psychosocial correlates: findings from a U.K. community study. *Epilepsia* 1996;37: 708–15.
- Tellez-Zenteno JF, Patten SB, Jette N, Williams J, Wiebe S. Psychiatric comorbidity in epilepsy: a population-based analysis. *Epilepsia* 2007;48:2336–44.
- Gilliam FG, Santos J, Vahle V, Carter J, Brown K, Hecimovic H. Depression in epilepsy: ignoring clinical expression of neuronal network dysfunction? *Epilepsia* 2004;45 (Suppl. 2):28–33.
- Hesdorffer DC, Ishihara L, Mynepalli L, Webb DJ, Weil J, Hauser WA. Epilepsy, suicidality, and psychiatric disorders: a bidirectional association. *Ann Neurol* 2012; 72:184–91.
- Adelow C, Andersson T, Ahlbom A, Tomson T. Hospitalization for psychiatric disorders before and after onset of unprovoked seizures/epilepsy. *Neurology* 2012;78: 396–401.
- Forsgren L, Nystrom L. An incident case-referent study of epileptic seizures in adults. *Epilepsy Res* 1990;6:66–81.
- Hesdorffer DC, Hauser WA, Annegers JF, Cascino G. Major depression is a risk factor for seizures in older adults. *Ann Neurol* 2000;47:246–9.
- Hesdorffer DC, Hauser WA, Olafsson E, Ludvigsson P, Kjartansson O. Depression and suicide attempt as risk factors for incident unprovoked seizures. *Ann Neurol* 2006; 59:35–41.
- Brandt C, Mula M. Anxiety disorders in people with epilepsy. *Epilepsy Behav* 2016; 59:87–91.
- Kobau R, Gilliam F, Thurman DJ. Prevalence of self-reported epilepsy or seizure disorder and its associations with self-reported depression and anxiety: results from the 2004 HealthStyles Survey. *Epilepsia* 2006;47:1915–21.
- Ottman R, Lipton RB, Ettinger AB, Cramer JA, Reed ML, Morrison A, et al. Comorbidities of epilepsy: results from the Epilepsy Comorbidities and Health (EPIC) survey. *Epilepsia* 2011;52:308–15.
- Rai D, Kerr MP, McManus S, Jordanova V, Lewis G, Brugha TS. Epilepsy and psychiatric comorbidity: a nationally representative population-based study. *Epilepsia* 2012; 53:1095–103.
- Ettinger AB, Copeland LA, Zeber JE, Van Cott AC, Pugh MJ. Are psychiatric disorders independent risk factors for new-onset epilepsy in older individuals? *Epilepsy Behav* 2010;17:70–4.
- Kanner AM, Rivas-Grajales AM. Psychosis of epilepsy: a multifaceted neuropsychiatric disorder. *CNS Spectr* 2016;21:247–57.
- Adachi N, Ito M, Kanemoto K, Akanuma N, Okazaki M, Ishida S, et al. Duration of postictal psychotic episodes. *Epilepsia* 2007;48:1531–7.
- Chen Z, Lusicic A, O'Brien TJ, Velakoulis D, Adams SJ, Kwan P. Psychotic disorders induced by antiepileptic drugs in people with epilepsy. *Brain* 2016;139: 2668–78.
- Adachi N, Matsuura M, Okubo Y, Oana Y, Takei N, Kato M, et al. Predictive variables of interictal psychosis in epilepsy. *Neurology* 2000;55:1310–4.
- Qin P, Xu H, Laursen TM, Vestergaard M, Mortensen PB. Risk for schizophrenia and schizophrenia-like psychosis among patients with epilepsy: population based cohort study. *BMJ* 2005;331:23.
- Mendez MF, Grau R, Doss RC, Taylor JL. Schizophrenia in epilepsy: seizure and psychosis variables. *Neurology* 1993;43:1073–7.
- Clancy MJ, Clarke MC, Connor DJ, Cannon M, Cotter DR. The prevalence of psychosis in epilepsy: a systematic review and meta-analysis. *BMC Psychiatry* 2014;14:75.
- Kanemoto K, Takeuchi J, Kawasaki J, Kawai I. Characteristics of temporal lobe epilepsy with mesial temporal sclerosis, with special reference to psychotic episodes. *Neurology* 1996;47:1199–203.
- Wotton CJ, Goldacre MJ. Coexistence of schizophrenia and epilepsy: record-linkage studies. *Epilepsia* 2012;53:e71–4.
- Chang YT, Chen PC, Tsai JJ, Sung FC, Chin ZN, Kuo HT, et al. Bidirectional relation between schizophrenia and epilepsy: a population-based retrospective cohort study. *Epilepsia* 2011;52:2036–42.
- Lin JJ, Mula M, Hermann BP. Uncovering the neurobehavioural comorbidities of epilepsy over the lifespan. *Lancet* 2012;380:1180–92.
- Meador KJ. Cognitive and memory effects of the new antiepileptic drugs. *Epilepsy Res* 2006;68:63–7.
- Mula M, Trimble MR. Antiepileptic drug-induced cognitive adverse effects: potential mechanisms and contributing factors. *CNS Drugs* 2009;23:121–37.
- Mula M, von Oertzen TJ, Cock HR, Lozsadi DA, Agrawal N. Clinical correlates of memory complaints during AED treatment. *Acta Neurol Scand* 2016;134:368–73.
- Marino SE, Meador KJ, Loring DW, Okun MS, Fernandez HH, Fessler AJ, et al. Subjective perception of cognition is related to mood and not performance. *Epilepsy Behav* 2009;14:459–64.
- Helmstaedter C, Elger CE. Chronic temporal lobe epilepsy: a neurodevelopmental or progressively dementing disease? *Brain* 2009;132:2822–30.

- [34] Taylor J, Kolamunnage-Dona R, Marson AG, Smith PE, Aldenkamp AP, Baker GA, et al. Patients with epilepsy: cognitively compromised before the start of antiepileptic drug treatment? *Epilepsia* 2010;51:48–56.
- [35] Ravizza T, Onat FY, Brooks-Kayal AR, Depaulis A, Galanopoulou AS, Mazarati A, et al. WONOEP appraisal: biomarkers of epilepsy-associated comorbidities. *Epilepsia* 2017;58:331–42.
- [36] Boylan LS, Flint LA, Labovitz DL, Jackson SC, Starner K, Devinsky O. Depression but not seizure frequency predicts quality of life in treatment-resistant epilepsy. *Neurology* 2004;62:258–61.
- [37] Hitiris N, Mohanraj R, Norrie J, Sills GJ, Brodie MJ. Predictors of pharmacoresistant epilepsy. *Epilepsy Res* 2007;75:192–6.
- [38] Nogueira MH, Yasuda CL, Coan AC, Kanner AM, Cendes F. Concurrent mood and anxiety disorders are associated with pharmacoresistant seizures in patients with MTLE. *Epilepsia* 2017;58:1268–76.
- [39] Kanner AM, Byrne R, Chicharro A, Wu J, Frey M. A lifetime psychiatric history predicts a worse seizure outcome following temporal lobectomy. *Neurology* 2009;72:793–9.
- [40] Koch-Stoecker SC, Bien CG, Schulz R, May TW. Psychiatric lifetime diagnoses are associated with a reduced chance of seizure freedom after temporal lobe surgery. *Epilepsia* 2017;58:983–93.
- [41] Cramer JA, Blum D, Reed M, Fanning K. The influence of comorbid depression on seizure severity. *Epilepsia* 2003;44:1578–84.
- [42] Gur-Ozmen S, Mula M, Agrawal N, Cock HR, Lozsadi D, von Oertzen TJ. The effect of depression and side effects of antiepileptic drugs on injuries in patients with epilepsy. *Eur J Neurol* 2017;24:1135–9.
- [43] Fazel S, Wolf A, Langstrom N, Newton CR, Lichtenstein P. Premature mortality in epilepsy and the role of psychiatric comorbidity: a total population study. *Lancet* 2013;382:1646–54.
- [44] Aaberg KM, Bakken IJ, Lossius MI, Lund Soraas C, Haberg SE, Stoltenberg C, et al. Comorbidity and childhood epilepsy: a nationwide registry study. *Pediatrics* 2016;138.
- [45] Oh A, Thurman DJ, Kim H. Comorbidities and risk factors associated with newly diagnosed epilepsy in the U.S. pediatric population. *Epilepsy Behav* 2017;75:230–6.
- [46] Brodie MJ, Besag F, Ettinger AB, Mula M, Gobbi G, Comai S, et al. Epilepsy, antiepileptic drugs, and aggression: an evidence-based review. *Pharmacol Rev* 2016;68:563–602.
- [47] Salpekar JA, Mishra G, Hauptman AJ. Key issues in addressing the comorbidity of depression and pediatric epilepsy. *Epilepsy Behav* 2015;46:12–8.
- [48] Guilfoyle SM, Follansbee-Junger K, Smith AW, Combs A, Ollier S, Hater B, et al. Antiepileptic drug behavioral side effects and baseline hyperactivity in children and adolescents with new onset epilepsy. *Epilepsia* 2018;59:146–54.
- [49] Davico C, Canavese C, Vittorini R, Gandione M, Vitiello B. Anticonvulsants for psychiatric disorders in children and adolescents: a systematic review of their efficacy. *Front Psychol* 2018;9:270.
- [50] Caplan R, Siddarth P, Stahl L, Lanphier E, Vona P, Gurbani S, et al. Childhood absence epilepsy: behavioral, cognitive, and linguistic comorbidities. *Epilepsia* 2008;49:1838–46.
- [51] Salpekar JA, Mishra G. Key issues in addressing the comorbidity of attention deficit hyperactivity disorder and pediatric epilepsy. *Epilepsy Behav* 2014;37:310–5.
- [52] Lee HJ, Kim EH, Yum MS, Ko TS, Kim HW. Attention profiles in childhood absence epilepsy compared with attention-deficit/hyperactivity disorder. *Brain and Development* 2018;40:94–9.
- [53] Verrotti A, Matricardi S, Rinaldi VE, Prezioso G, Coppola G. Neuropsychological impairment in childhood absence epilepsy: review of the literature. *J Neurol Sci* 2015;359:59–66.
- [54] Li Q, Cao W, Liao X, Chen Z, Yang T, Gong Q, et al. Altered resting state functional network connectivity in children absence epilepsy. *J Neurol Sci* 2015;354:79–85.
- [55] Vega C, Guo J, Killory B, Danielson N, Vestal M, Berman R, et al. Symptoms of anxiety and depression in childhood absence epilepsy. *Epilepsia* 2011;52:e70–4.
- [56] Scharfman HE, Kanner AM, Friedman A, Blumcke I, Crocker CE, Cendes F, et al. Epilepsy as a network disorder (2): what can we learn from other network disorders such as dementia and schizophrenia, and what are the implications for translational research? *Epilepsy Behav* 2017;78:302–12.
- [57] Reilly C, Agnew R, Neville BG. Depression and anxiety in childhood epilepsy: a review. *Seizure* 2011;20:589–97.
- [58] Maryam S, Parviz B. Depression in children and adolescents with epilepsy: a 15 year research review of prevalence, and demographic and seizure related correlates. *Iran J Pediatr* 2013;23:1–7.
- [59] Han SH, Lee SA, Eom S, Kim HD, Korean Qo LiESG. Family factors contributing to emotional and behavioral problems in Korean adolescents with epilepsy. *Epilepsy Behav* 2016;56:66–72.
- [60] Liu X, Han Q. Depression and anxiety in children with benign childhood epilepsy with centrotemporal spikes (BCECTS). *BMC Pediatr* 2016;16:128.
- [61] Roeder R, Roeder K, Asano E, Chugani HT. Depression and mental health help-seeking behaviors in a predominantly African American population of children and adolescents with epilepsy. *Epilepsia* 2009;50:1943–52.
- [62] Coyle JT, Pine DS, Charney DS, Lewis L, Nemeroff CB, Carlson GA, et al. Depression and bipolar support alliance consensus statement on the unmet needs in diagnosis and treatment of mood disorders in children and adolescents. *J Am Acad Child Adolesc Psychiatry* 2003;42:1494–503.
- [63] Dunn DW, Besag F, Caplan R, Aldenkamp A, Gobbi G, Sillanpaa M. Psychiatric and behavioural disorders in children with epilepsy (ILAE Task Force Report): anxiety, depression and childhood epilepsy. *Epileptic Disord* 2016;18(Suppl1):S24–30.
- [64] Vona P, Siddarth P, Sankar R, Caplan R. Obstacles to mental health care in pediatric epilepsy: insight from parents. *Epilepsy Behav* 2009;14:360–6.
- [65] Hamid H, Blackmon K, Cong X, Dziura J, Atlas LY, Vickrey BG, et al. Mood, anxiety, and incomplete seizure control affect quality of life after epilepsy surgery. *Neurology* 2014;82:887–94.
- [66] Puka K, Smith ML. Predictors of long-term quality of life after pediatric epilepsy surgery. *Epilepsia* 2015;56:873–81.
- [67] Salpekar JA, Berl MM, Havens K, Cushner-Weinstein S, Conry JA, Pearl PL, et al. Psychiatric symptoms in children prior to epilepsy surgery differ according to suspected seizure focus. *Epilepsia* 2013;54:1074–82.
- [68] Pereira A, Valente KD. Severity of depressive symptomatology and functional impairment in children and adolescents with temporal lobe epilepsy. *Seizure* 2013;22:708–12.
- [69] Caetano SC, Fonseca M, Hatch JP, Olvera RL, Nicoletti M, Hunter K, et al. Medial temporal lobe abnormalities in pediatric unipolar depression. *Neurosci Lett* 2007;427:142–7.
- [70] Besag F, Gobbi G, Caplan R, Sillanpaa M, Aldenkamp A, Dunn DW. Psychiatric and behavioural disorders in children with epilepsy (ILAE Task Force Report): epilepsy and ADHD. *Epileptic Disord* 2016;18(Suppl1):S8–S15.
- [71] Cohen R, Senecy Y, Shuper A, Inbar D, Chodick G, Shalev V, et al. Prevalence of epilepsy and attention-deficit hyperactivity (ADHD) disorder: a population-based study. *J Child Neurol* 2013;28:120–3.
- [72] Torres AR, Whitney J, Gonzalez-Heydrich J. Attention-deficit/hyperactivity disorder in pediatric patients with epilepsy: review of pharmacological treatment. *Epilepsy Behav* 2008;12:217–33.
- [73] Ekinci O, Okuyaz C, Erdogan S, Gunes S, Ekinci N, Kalinli M, et al. Attention-deficit hyperactivity disorder (ADHD) in epilepsy and primary ADHD: differences in symptom dimensions and quality of life. *J Child Neurol* 2017;32:1083–91.
- [74] Gascoigne MB, Smith ML, Barton B, Webster R, Gill D, Lah S. Attention deficits in children with epilepsy: preliminary findings. *Epilepsy Behav* 2017;67:7–12.
- [75] Zhang DQ, Li FH, Zhu XB, Sun RP. Clinical observations on attention-deficit hyperactivity disorder (ADHD) in children with frontal lobe epilepsy. *J Child Neurol* 2012;24:54–7.
- [76] Silvestri R, Gagliano A, Calarese T, Arico I, Cedro C, Condurso R, et al. Ictal and interictal EEG abnormalities in ADHD children recorded over night by video-polysomnography. *Epilepsy Res* 2007;75:130–7.
- [77] Fonseca LC, Tedrus GM, Moraes C, Vicente Machado A, Almeida MP, Oliveira DO. Epileptiform abnormalities and quantitative EEG in children with attention-deficit/hyperactivity disorder. *Arq Neuropsiquiatr* 2008;66:462–7.
- [78] Kanemura H, Sano F, Tando T, Hosaka H, Sugita K, Aihara M. EEG improvements with antiepileptic drug treatment can show a high correlation with behavioral recovery in children with ADHD. *Epilepsy Behav* 2013;27:443–8.
- [79] Chou IC, Chang YT, Chin ZN, Muo CH, Sung FC, Kuo HT, et al. Correlation between epilepsy and attention deficit hyperactivity disorder: a population-based cohort study. *PLoS One* 2013;8:e57926.
- [80] Sillanpaa M, Besag F, Aldenkamp A, Caplan R, Dunn DW, Gobbi G. Psychiatric and behavioural disorders in children with epilepsy (ILAE Task Force Report): epidemiology of psychiatric/behavioural disorder in children with epilepsy. *Epileptic Disord* 2016;18(Suppl1):S2–7.
- [81] Ekinci O, Arman AR, Isik U, Bez Y, Berkem M. EEG abnormalities and epilepsy in autistic spectrum disorders: clinical and familial correlates. *Epilepsy Behav* 2010;17:178–82.
- [82] Wheless JW, Simos PG, Butler JJ. Language dysfunction in epileptic conditions. *Semin Pediatr Neurol* 2002;9:218–28.
- [83] Trauner DA. Behavioral correlates of epileptiform abnormalities in autism. *Epilepsy Behav* 2015;47:163–6.
- [84] Jacob J. Cortical interneuron dysfunction in epilepsy associated with autism spectrum disorders. *Epilepsia* 2016;57:182–93.
- [85] Salpekar J. Neuropsychiatric effects of epilepsy in developmental disorders. *Curr Opin Psychiatry* 2018;31:109–15.